

**Table 1. Summary<sup>a</sup> of Range of Differences in Second Doses of Diazepam Nasal Spray Between Treatment Period 1 and Treatment Period 2 and Mean Number of Seizure Cluster Events by Period, Based on 2-Month Cutoffs and Number of Events**

<b>Cutoff: Months per Period</b>	<b>Cutoffs: Events per Period<sup>b</sup></b>	<b>No. of Subjects Based on Cutoff<sup>c</sup></b>	<b>Difference (Period 2 minus Period 1)</b>	<b>Mean No. of Events, Period 1</b>	<b>Mean No. of Events, Period 2</b>	<b>Wilcoxon Signed Rank Test <i>P</i> value</b>
2	1–7	12–101	2.9% to 8.7%	4.8–9.8	4.6–10.3	0.021–0.214
3	1–10	8–103	–1.8% to 12.0%	7.3–14.1	6.0–13.6	0.063–0.805
4	1–13	9–99	–0.9% to 3.8%	8.8–19.0	8.3–18.1	0.432–0.892
5	1–18	8–91	–4.5% to –0.1%	11.0–24.4	10.0–23.1	0.034–0.703
6	1–18	9–62	–3.6% to 2.9%	14.0–26.1	11.4–26.0	0.129–0.880
7	1–18	8–43	–3.8% to –0.9%	16.1–27.6	13.0–29.0	0.074–0.547
8	1–17	8–36	–2.0% to –0.5%	17.1–29.0	13.4–31.0	0.173–0.855
9	1–18	8–32	–4.7% to –1.3%	19.5–32.5	15.8–34.4	0.049–0.557
10	1–18	8–24	–2.2% to 1.4%	23.8–35.9	18.4–37.9	0.268–1.000
11	1–12	8–13	–1.0% to 1.8%	29.2–35.9	23.0–32.9	0.426–1.000
12	1–12	8–12	–1.5% to 1.1%	32.8–37.4	25.0–34.4	0.426–0.844
13	1–14	8–12	–0.5% to 2.3%	36.0–41.6	27.1–37.9	0.652–0.938
14	1–7	8–10	–2.6% to –2.3%	37.5–39.5	34.2–41.8	0.250–0.375
15	1–9	8–10	–2.0% to –1.8%	39.7–42.1	35.7–43.3	0.383–0.469

<sup>a</sup>Summary of all 191 analyses.

<sup>b</sup>Reflects number of individual analyses summarized in the row (eg, 2 month row represents 7 individual analyses).  
Minimum 8 subjects.